



Development of immune response and decay of maternal immunity after vaccination with C-strain

Rangelova, Desislava Yordanova; Nielsen, Jens; Uttenthal, Åse

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Rangelova, D. Y., Nielsen, J., & Uttenthal, Å. (2011). *Development of immune response and decay of maternal immunity after vaccination with C-strain*. Abstract from 8th ESVV International Pestivirus Symposium, Hanover, Lower Saxony, Germany. <http://www.pestivirus2011.de/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



8th ESVV Pestivirus Symposium
September 25 – 28, 2011
Hannover, Germany

Session 3: Virus-host interactions - poster ID 51

Development of immune response and decay of maternal immunity after vaccination with C-strain

Desislava Rangelova, Jens Nielsen, Åse Uttenthal.

Technical University of Denmark, National Veterinary Institute, Lindholm, Denmark

The most widely used vaccine towards CSFV is the C strain, modified live vaccine. The aims of the present study were to evaluate the humoral immune response after vaccination with C-strain and the impact of maternal immunity in piglets obtained from C-strain vaccinated sows.

Two experiments were conducted.

Experiment A involved 12, 5-weeks-piglets vaccinated intramuscularly. Antibodies were measured twice weekly up to 28 days post vaccination.

Experiment B followed the decay of maternal immunity in piglets obtained from four sows vaccinated intramuscularly at 86 days of gestation. The serological response in the sows was measured 14 days post vaccination, at farrowing, two and four weeks post farrowing. In the outcome piglets the antibody level was measured weekly up to 35 days of age.

Serum samples were tested by neutralization and by a routine blocking CSFV ELISA, respectively. The results obtained with the first experiment showed that all pigs seroconverted two weeks post vaccination. The antibody level increased during the first four weeks post vaccination. Also the four vaccinated sows were antibody positive two weeks post vaccination. Negative precolostral samples from the pigs showed that that transplacental transmission of antibodies had not occurred. After uptake of colostrums, however, the piglets seroconverted. The duration of maternal immunity in pigs obtained from the vaccinated sows is in progress and the results will be presented at the Symposium.

Acknowledgement: The study is part of CSFVgoDIVA (EU-227003).